


ref. 1



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 211 243 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication: **05.06.2002 Bulletin 2002/23**
(51) Int Cl.7: **C07D 211/32, A61K 31/445, A61P 25/28**
(21) Application number: **02005248.6**
(22) Date of filing: **06.06.1997**

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE

(30) Priority: **07.06.1996 JP 14629396**
27.12.1996 WOPCT/JP96/03881
30.12.1996 US 774802

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
97924337.5 / 1 019 374

(71) Applicant: **Eisai Co., Ltd.**
Tokyo 112-0002 (JP)

(72) Inventors:
• **Watanabe, Hideaki**
Ushiku-shi, Ibaraki 300-12 (JP)
• **Imai, Akio**
Kashima-gun, Ibaraki 314-02 (JP)

- **Kajima, Takashi**
Ryugasaki-shi, Ibaraki 301 (JP)
- **Ishihama, Yasushi**
Kitasoma-gun, Ibaraki 302-0124 (JP)
- **Ohtsuka, Akiyo**
Tsukuba-shi, Ibaraki 305 (JP)
- **Tanaka, Tomohide**
Kashima-gun, Ibaraki 314-02 (JP)
- **Nabaru, Yukio**
Kashima-gun, Ibaraki 314-04 (JP)

(74) Representative: **HOFFMANN - EITLE**
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

Remarks:

This application was filed on 08 - 03 - 2002 as a divisional application to the application mentioned under INID code 62.

(54) **Polymorphs of donepezil hydrochloride and process for production**

(57) Donepezil hydrochloride, 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride, is provided here in the form of four polymorphs which are stable against heat and humidity in the pharmaceutical use. They can be industrially pro-

duced. They are specified by peaks in X-ray powder diffraction pattern and absorption peaks in infrared absorption spectra in potassium bromide.

EP 1 211 243 A1

Description

[Field of the invention]

5 **[0001]** The present invention relates to the stable polymorphs of Donepezil hydrochloride, that is, 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methyl-piperidine hydrochloride, disclosed in the example 4 of US-A 4,895,841 or EP-A 296560, having an excellent efficacy as pharmaceuticals, and industrial processes for producing them.

[Background of the invention]


10 **[0002]** Donepezil hydrochloride shows the acetylcholine esterase-inhibitory action and is useful for the treatment of all kinds of senile dementia, in particular being useful for prevention, treatment and amelioration of Alzheimer Disease. Donepezil hydrochloride is administered orally as usual and it may be placed for distribution and storage in a period of time before the administration. It may then be stored at patient's home for about one month at the maximum because
 15 of the property of the target disease. The stability of this medicinal substance (bulk pharmaceutical chemicals) against heat and humidity during the storage period is very important. A more stable medicinal substance of Donepezil hydrochloride is, therefore, desired. It is not known, however, that polymorphs of Donepezil hydrochloride exist. No sufficiently stable medicinal substance of Donepezil hydrochloride has been found.

[Prior Arts]

20 **[0003]** US-A 4,895,841 discloses in Example 4 that recrystallization of the crude product mixture of Donepezil hydrochloride from ethanol/isopropyl ether afforded a purified Donepezil hydrochloride. If there is a more stable crystalline form of Donepezil hydrochloride for a long period, it is more practical for distribution and storage.

[Summary of the invention]

25 **[0004]** Regarding the foregoing problems, the present inventors have proceeded with extensive research. As a result, it has been found that novel polymorphs of Donepezil hydrochloride (I) to (IV) and (V) can be produced and have an excellent stability, establishing the present invention. The present invention offers the five forms or species of novel polymorphs of Donepezil hydrochloride and industrially excellent processes for producing them.
 30 In detail, the present invention relates to the five polymorphs (I) to (V) of Donepezil hydrochloride represented by the following chemical structure, the polymorphs being specified by the peaks appearing in the powder X-ray diffraction pattern and infrared absorption spectra in potassium bromide.



(Method, and condition of the measurement of X-ray diffraction patterns)

(1) Method of the measurement

45 **[0005]** X-ray diffraction patterns were measured on each 100mg of the samples by the following condition.

(2) Condition of the measurement

[0006]

55

| | |
|---------|----------|
| Target | Cu |
| Filter | monochro |
| Voltage | 40 KV |
| Current | 20 mA |

EP 1 211 243 A1

(continued)

| | |
|-------------|---------------------|
| Slit | DS 1, RS 0.15, SS 1 |
| Scan speed | 2 deg/min. |
| Range | 5-30 |
| Step/Sample | 0.02 deg |

(Method and condition of the measurement of infrared absorption)

[0007] Infrared absorption spectra in potassium bromide were measured according to the general method recorded in the Japanese Pharmacopoeia.

(1) Polymorphs (I)

[0008] Peaks in the powder X-ray diffraction pattern are:

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 9.94 | 24 |
| 10.60 | 19 |
| 12.66 | 69 |
| 13.12 | 55 |
| 13.66 | 44 |
| 13.86 | 40 |
| 14.92 | 49 |
| 15.26 | 17 |
| 16.08 | 35 |
| 16.86 | 34 |
| 17.50 | 34 |
| 17.58 | 42 |
| 18.42 | 20 |
| 19.28 | 27 |
| 19.80 | 45 |
| 19.94 | 45 |
| 21.22 | 100 |
| 22.00 | 32 |
| 22.54 | 31 |
| 22.98 | 49 |
| 23.60 | 56 |
| 23.78 | 75 |
| 23.92 | 78 |
| 26.46 | 33 |
| 28.02 | 25 |
| 29.50 | 37 |

[0009] Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

463, 502, 563, 589, 604, 701, 750, 759, 799, 860, 922, 947, 972, 1012, 1038, 1104, 1120, 1128, 1175, 1192, 1218, 1250, 1267, 1316, 1368, 1410, 1433, 1440, 1455, 1472, 1502, 1591, 1606, 1644, 1684, 2412, 2530, 2559, 2595, 2620, 2717, 2840, 2858, 2924, 3004, 3074, 3259, 3373, 3547, 3589 cm^{-1} .

(2) Polymorphs (II)

[0010] Peaks in the powder X-ray diffraction pattern are:

EP 1 211 243 A1

| | Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|----|---|--------------------------|
| 5 | 7.40 | 8 |
| | 9.88 | 100 |
| | 12.36 | 13 |
| | 15.54 | 40 |
| | 16.10 | 38 |
| 10 | 16.22 | 38 |
| | 16.48 | 35 |
| | 17.30 | 17 |
| | 18.04 | 20 |
| 15 | 18.44 | 17 |
| | 18.84 | 19 |
| | 19.34 | 19 |
| | 19.84 | 47 |
| | 21.16 | 24 |
| 20 | 22.40 | 19 |
| | 23.18 | 33 |
| | 24.02 | 22 |
| | 24.92 | 25 |
| 25 | 25.72 | 27 |
| | 26.40 | 18 |
| | 27.22 | 14 |

[0011] Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

699, 748, 762, 845, 947, 1009, 1035, 1067, 1103, 1118, 1129, 1174, 1193, 1206, 1222, 1247, 1267, 1317, 1365, 1422, 1436, 1456, 1465, 1502, 1592, 1607, 1688, 2412, 2489, 2627, 2846, 2868, 2913, 2928, 3435 cm^{-1} .

(3) Polymorphs (III)

[0012] Peaks in the powder X-ray diffraction pattern are:

| | Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|----|---|--------------------------|
| 40 | 6.56 | 30 |
| | 9.94 | 8 |
| | 13.00 | 17 |
| | 15.00 | 47 |
| 45 | 15.26 | 14 |
| | 15.74 | 6 |
| | 16.48 | 35 |
| | 17.42 | 4 |
| | 18.10 | 21 |
| 50 | 18.50 | 56 |
| | 19.50 | 17 |
| | 20.10 | 32 |
| | 20.94 | 21 |
| 55 | 21.66 | 100 |
| | 22.32 | 25 |
| | 22.92 | 17 |

EP 1 211 243 A1

(continued)

| Diffraction angles ($2\theta,^\circ$) | Intensity (I/I_0) |
|--|--------------------------|
| 23.92 | 19 |
| 24.68 | 17 |
| 26.00 | 44 |
| 27.20 | 23 |
| 28.02 | 29 |
| 28.22 | 40 |
| 28.60 | 13. |

[0013] Wave numbers (cm^{-1}) of infrared absorption spectrum in potassium bromide are:

559, 641, 648, 702, 749, 765, 786, 807, 851, 872, 927, 949, 966, 975, 982, 1007, 1034, 1071, 1080, 1111, 1119, 1131, 1177, 1190, 1205, 1217, 1230, 1250, 1265, 1292-, 1313, 1367, 1389, 1420, 1438, 1453, 1461, 1470, 1500, 1589, 1605, 1697, 2407, 2419, 2461, 2624, 2641, 2651, 2667, 2837, 2848, 2873, 2924, 2954, 2961, 2993, 3007, 3377, 3433 cm^{-1} .

(4) Polymorphs (IV)

[0014] Peaks in the powder X-ray diffraction pattern are:

| Diffraction angles ($2\theta,^\circ$) | Intensity (I/I_0) |
|--|--------------------------|
| 6.24 | 15 |
| 9.66 | 12 |
| 11.04 | 22 |
| 12.12 | 24 |
| 12.54 | 67 |
| 12.76 | 61 |
| 13.98 | 27 |
| 14.42 | 15 |
| 14.88 | 11 |
| 16.34 | 12 |
| 17.46 | 100 |
| 18.12 | 25 |
| 18.60 | 32 |
| 19.06 | 15 |
| 19.98 | 74 |
| 20.42 | 41 |
| 20.62 | 34 |
| 21.30 | 48 |
| 21.80 | 63 |
| 22.58 | 78 |
| 23.04 | 46 |
| 24.00 | 32 |
| 24.54 | 49 |
| 25.14 | 90 |
| 25.36 | 99 |
| 26.06 | 34 |
| 28.10 | 41 |
| 28.58 | 39 |

EP 1 211 243 A1

(continued)

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 29.30 | 31 |
| 29.44 | 28. |

[0015] Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

401, 431, 459, 467, 490, 506, 518, 561, 586, 606, 631, 651, 709, 758, 766, 857, 944, 1009, 1041, 1106, 1119, 1132, 1213, 1225, 1265, 1304, 1318, 1429, 1458, 1470, 1500, 1589, 1605, 1630, 1647, 1683, 2562, 2577, 2608, 2634, 2689, 2717, 2836, 2924, 2949, 2989, 3007, 3032, 3061, 3322, 3376, 3422 cm^{-1} .

[0016] Additional analysis data of polymorphs (I) to (V) are represented as following.

(1) Polymorphs (I)

[0017] Peaks in the powder X-ray diffraction pattern are:

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 9.82 | 13 |
| 10.48 | 18 |
| 12.52 | 93 |
| 13.02 | 69 |
| 13.52 | 34 |
| 13.74 | 37 |
| 14.78 | 31 |
| 16.00 | 45 |
| 16.76 | 37 |
| 17.46 | 34 |
| 19.18 | 26 |
| 19.66 | 32 |
| 21.04 | 100 |
| 21.16 | 82 |
| 22.92 | 52 |
| 23.82 | 72 |
| 24.14 | 32. |

[0018] Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

562.7, 603.2, 700.4, 749.6, 798.1, 859.2, 896.0, 921.3, 946.3, 971.8, 1038.0, 1119.3, 1216.8, 1266.0, 1315.4, 1367.7, 1454.1, 1501.5, 1537.8, 1555.9, 1590.7, 1643.7, 1681.9, 2350.9, 2534.0, 2922.1, 3381.8, 3585.2 cm^{-1} .

(2) Polymorphs (II)

[0019] Peaks in the powder X-ray diffraction pattern are:

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 10.10 | 76 |
| 12.64 | 14 |
| 15.74 | 85 |

EP 1 211 243 A1

(continued)

5

10

15

20

25

30

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 15.82 | 86 |
| 16.20 | 100 |
| 16.46 | 87 |
| 17.40 | 50 |
| 17.50 | 48 |
| 17.88 | 31 |
| 18.36 | 28 |
| 18.58 | 51 |
| 18.66 | 46 |
| 19.48 | 42 |
| 20.18 | 81 |
| 20.80 | 36 |
| 22.26 | 45 |
| 23.38 | 86 |
| 23.52 | 59 |
| 24.06 | 34 |
| 24.32 | 55 |
| 25.14 | 44 |
| 25.44 | 50 |
| 25.72 | 39 |
| 25.96 | 35 |
| 26.14 | 25 |
| 28.06 | 25 |
| 28.20 | 34 |
| 28.38 | 34 |

[0020] Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

35

560.1, 698.9, 749.1, 846.2, 947.6, 1036.1, 1119.3, 1222.7, 1266.4, 1318.7, 1364.1, 1458.3, 1500.9, 1522.3, 1534.0, 1542.6, 1560.2, 1570.3, 1592.0, 1637.0, 1647.9, 1654.4, 1689.5, 1718.3, 1734.7, 1751.7, 1773.9, 1793.8, 1830.7, 1846.0, 1870.1, 2345.1, 2489.9, 2927.9, 3448.1 cm^{-1} .

40

(3) Polymorphs (III)

[0021] Peaks in the powder X-ray diffraction pattern are:

45

50

55

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 6.48 | 21 |
| 9.84 | 7 |
| 12.96 | 19 |
| 14.94 | 45 |
| 15.20 | 13 |
| 16.44 | 31 |
| 18.04 | 20 |
| 18.46 | 55 |
| 19.44 | 17 |
| 20.02 | 30 |
| 20.86 | 20 |

EP 1 211 243 A1

(continued)

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 21.02 | 13 |
| 21.58 | 100 |
| 22.22 | 23 |
| 22.90 | 15 |
| 23.92 | 13 |
| 24.64 | 15 |
| 25.92 | 40 |
| 26.18 | 17 |
| 27.14 | 21 |
| 28.14 | 37 |
| 28.56 | 11 |
| 29.94 | 12. |

[0022] Wave numbers (cm^{-1}) of infrared absorption spectrum in potassium bromide are:

558.3, 641.1, 702.4, 748.5, 765.0, 786.1, 807.3, 850.8, 872.0, 926.8, 974.9, 1034.1, 1071.5, 1111.6, 1190.1, 1216.6, 1265.4, 1291.9, 1312.9, 1364.4, 1420.2, 1438.1, 1458.8, 1499.1, 1522.2, 1542.6, 1560.1, 1570.2, 1589.1, 1638.8, 1647.8, 1654.3, 1697.3, 1718.1, 1734.5, 1751.4, 1773.7, 1793.5, 1845.8, 2345.3, 2461.6, 2924.2, 3447.9 cm^{-1} .

(4) Polymorphs (IV)

[0023] Peaks in the powder X-ray diffraction pattern are:

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 9.64 | 11 |
| 10.92 | 11 |
| 12.46 | 63 |
| 12.72 | 17 |
| 13.86 | 27 |
| 14.42 | 12 |
| 17.36 | 100 |
| 18.54 | 39 |
| 19.90 | 37 |
| 21.18 | 35 |
| 21.74 | 39 |
| 22.48 | 60 |
| 22.96 | 36 |
| 24.10 | 17 |
| 25.28 | 70 |
| 28.00 | 27 |
| 28.50 | 27. |

[0024] Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

561.5, 709.0, 766.2, 786.3, 804.9, 857.0, 944.3, 979.3, 1041.5, 1118.7, 1264.6, 1318.7, 1364.1, 1458.1, 1499.2, 1542.5, 1560.1, 1588.1, 1636.6, 1647.8, 1654.3, 1684.3, 1718.2, 1734.4, 1751.4, 1773.7, 1793.5, 1830.5, 1845.8, 1870.1, 2344.8, 2369.3, 2719.2, 2922.9, 3324.0 cm^{-1} .

(5) Polymorphs (V)

[0025]

| | Diffraction angles (2 θ , °) | Intensity (I/I ₀) |
|----|--|----------------------------------|
| | 6.58 | 7 |
| | 6.86 | 27 |
| 5 | 10.12 | 32 |
| | 12.54 | 33 |
| 10 | 12.90 | 43 |
| | 13.64 | 64 |
| | 15.58 | 27 |
| 15 | 17.22 | 69 |
| | 18.44 | 72 |
| | 18.96 | 19 |
| | 19.30 | 25 |
| 20 | 19.64 | 19 |
| | 19.74 | 25 |
| | 20.30 | 19 |
| | 20.46 | 17 |
| 25 | 21.10 | 15 |
| | 21.96 | 100 |
| | 22.24 | 32 |
| | 24.22 | 63 |
| | 24.66 | 96 |
| 30 | 25.36 | 60 |
| | 26.14 | 15 |
| | 26.82 | 44 |
| | 27.52 | 24 |
| | 27.96 | 15 |
| 35 | 28.20 | 49 |
| | 29.58 | 13 |
| | 29.66 | 17 |
| | 29.76 | 17 |

[0026] Wave numbers (cm⁻¹) of infrared absorption spectra in potassium bromide are:

506.5, 559.7, 594.4, 698.0, 740.8, 805.1, 861.9, 948.5, 972.1, 1039.9, 1120.8, 1220.7, 1264.8, 1314.6, 1364.1, 1458.0, 1499.5, 1542.5, 1560.2, 1592.1, 1692.9, 2500.1, 2924.2, 2998.9, 3422.1 cm⁻¹.

[0027] Melting points of the novel Polymorphs (I) to (V) disclosed in the present invention are different from that of example 4 in US-A-4,895,841.

[0028] A melting point in US-P-4,895,841 is 211-212° C (decomposition).

| | |
|---------------------------------------|------------------------------|
| Melting point of the polymorph (I) | 225-226° C (decomposition), |
| Melting point of the polymorph (II) | 224-226° C (decomposition), |
| Melting point of the polymorph (III) | 229-231° C (decomposition), |
| Melting point of the polymorph (IV) | 226-228° C (decomposition), |
| Melting point of the polymorph (V) | 218-220° C (decomposition), |
| [Melting point of the amorphous form] | 220-222° C (decomposition)]. |

[0029] Furthermore, the thermogravimetric and differential thermal analysis (TG-DTA) of the present polymorphs

measured under the following condition show different patterns from the prior art. It is noted accordingly that their crystalline forms are completely different from the prior art.

{Method and condition of the thermogravimetric and differential thermal analysis (TG-DTA)}

[0030] About 3 to 6 mg of Samples were taken and subjected to thermal analysis under the following condition.

| | |
|-------------|-------------------|
| Reference | empty |
| Scan speed | 5° C/min. |
| Sampling | 0.7 sec. |
| Upper limit | 300° C |
| Lower limit | Room temperature. |

[0031] Detailed processes for preparing the novel polymorphs are as follows. In these processes, "Donepezil" means a free base of the Donepezil hydrochloride, i.e., 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine.

(1) Processes for preparing the polymorph (I) are:

- (1-1) Recrystallization of Donepezil hydrochloride from methanol,
- (1-2) Dissolving Donepezil hydrochloride in methanol, followed by addition of diethyl ether or diisopropyl ether,
- (1-3) Dissolving Donepezil in methanol, followed by addition of hydrochloric acid or hydrogen chloride,
- (1-4) Dissolving Donepezil in ethanol, followed by addition of diisopropyl ether, and hydrochloric acid or hydrogen chloride successively; or
- (1-5) Dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride, and diisopropyl ether successively, then filtration of the crystals immediately after separation.

Process (1-5) is preferred. This process is illustrated in Example 7.

(2) Processes for preparing the polymorph (II) are:

- (2-1) Dissolving Donepezil hydrochloride in ethanol, followed by addition of diethyl ether or diisopropyl ether,
- (2-2) Dissolving Donepezil hydrochloride in ethanol, followed by addition of diisopropyl ether, then filtration of the crystals after 10 to 30 minutes from the separation,
- (2-3) Dissolving Donepezil and hydrochloric acid or hydrogen chloride in ethanol, followed by addition of diethyl ether,
- (2-4) Dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride, then concentration,
- (2-5) Dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride, and diisopropyl ether successively; or
- (2-6) Dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride, and diisopropyl ether successively, then filtration of the crystals after 10 to 60 minutes, preferably 10 to 30 minutes from the separation.

Process (2-6) is preferred. This process is illustrated in Example 14.

(3) Processes for preparing the polymorph (III) are:

- (3-1) Dissolving Donepezil hydrochloride in ethanol, followed by addition of diethyl ether,
- (3-2) Dissolving Donepezil hydrochloride in dichloromethane, followed by addition of n-hexane,
- (3-3) Dissolving Donepezil in acetone, followed by addition of hydrochloric acid or hydrogen chloride,
- (3-4) Dissolving Donepezil in ethyl acetate, followed by addition of hydrochloric acid or hydrogen chloride,
- (3-5) Dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride, then addition of at least one solvent selected from diethyl ether, diisopropyl ether and n-hexane,
- (3-6) According to a process (3-5) wherein the selected solvent is diisopropyl ether, filtration of the crystals after 1 hour, preferably 2 hours, more preferably 6 hours from the separation; or
- (3-7) Heating of the polymorph (I) or (II).

Processes (3-5) and (3-6) are preferred. These processes are illustrated, respectively, in Examples 23 and 18.

(4) A process for preparing the polymorph (IV) is:

(4-1) Humidification of the polymorph (II).

Process (4-1) is illustrated in Example 24.

[0032] The aforementioned processes (1-5), (2-6) and (3-6) comprise dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride, and then adding diisopropyl ether. Anyone of these processes can produce polymorph (I), (II) or (III) by controlling an interval in time taken from recrystallization to isolation of crystals by filtration. These time lag may differ by the crystallization conditions, such as temperature, stirring velocity and a volume of a solvent. The following may be usually used.

- (1) Filtration of the crystals immediately after separation affords the polymorph (I).
- (2) Filtration of the crystals after 10 to 60 minutes, preferably 10 to 30 minutes from the separation affords the polymorph (II).
- (3) Filtration of the crystals after 1 hour, preferably 2 hours, more preferably 6 hours from the separation affords the polymorph (III).

[0033] Further Detailed processes for preparing the novel polymorphs are as follows.

(1) Processes for preparing the polymorph (I) are:

- (1-6) Dissolving Donepezil in methanol, followed by addition of hydrochloric acid or hydrogen chloride,
- (1-7) Dissolving Donepezil in methanol, followed by addition of hydrochloric acid or hydrogen chloride and addition of tert-butyl methyl ether, diisopropyl ether or ethyl acetate successively,
- (1-8) Dissolving Donepezil in ethanol, tetrahydrofuran or acetonitrile, followed by addition of hydrochloric acid or hydrogen chloride,
- (1-9) Dissolving Donepezil hydrochloride in methanol, followed by addition of tert-butyl methyl ether, ethyl acetate or n-hexane,
- (1-10) Recrystallization of Donepezil hydrochloride from ethanol; or
- (1-11) Dissolving Donepezil hydrochloride in ethanol, followed by addition of tert-butyl methyl ether.

Process (1-7) is preferred. This process is illustrated in Example 30, 31 and 32.
Process (1-9) is also preferred. This process is illustrated in Example 39, 40 and 41.

(2) Processes for preparing the polymorph (II) are:

- (2-7) Dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride and addition of tert-butyl methyl ether successively,
- (2-8) Dissolving Donepezil in isopropyl alcohol, acetone or tetrahydrofuran, followed by addition of hydrochloric acid or hydrogen chloride,
- (2-9) Dissolving Donepezil in methylene chloride, followed by addition of hydrochloric acid or hydrogen chloride and addition of diisopropyl ether successively,
- (2-10) Dissolving Donepezil hydrochloride in ethanol, followed by addition of tert-butyl methyl ether or diisopropyl ether and stirring blow 10 °C,
- (2-11) Dissolving Donepezil hydrochloride in methylene chloride, followed by addition of tert-butyl methyl ether or diisopropyl ether; or
- (2-12) Heating the polymorph (I) or amorphous form of Donepezil hydrochloride.

Process (2-7) is preferred. This process is illustrated in Example 45.
Process (2-10) is also preferred. This process is illustrated in Example 52, 53 and 54.

(3) Processes for preparing the polymorph (III) are:

- (3-8) Dissolving Donepezil in methanol, followed by addition of hydrochloric acid or hydrogen chloride and addition of acetone successively,

- (3-9) Dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride and addition of tert-butyl methyl ether successively,
 (3-10) Dissolving Donepezil in acetonitrile, acetone, a mixture of acetone and water, tetrahydrofuran or N,N-dimethylformamide, followed by addition of hydrochloric acid or hydrogen chloride,
 5 (3-11) Dissolving Donepezil in ethyl acetate, followed by addition of hydrochloric acid or hydrogen chloride and addition of tert-butyl methyl ether successively,
 (3-12) Dissolving Donepezil in dimethylsulfoxide, followed by addition of hydrochloric acid or hydrogen chloride and addition of tert-butyl methyl ether successively,
 10 (3-13) Dissolving Donepezil in toluene, followed by addition of hydrochloric acid or hydrogen chloride,
 (3-14) Recrystallization of Donepezil hydrochloride from methanol not lower than 10°C,
 (3-15) Dissolving Donepezil hydrochloride in methanol, followed by addition of tert-butyl methyl ether or acetonitrile,
 (3-16) Dissolving Donepezil hydrochloride in ethanol, followed by addition of tert-butyl methyl ether or acetonitrile and stirring not lower than 10°C,
 15 (3-17) Dissolving Donepezil hydrochloride in N,N-dimethylformamide or dimethylsulfoxide, followed by addition of tert-butyl methyl ether,
 (3-18) Recrystallization of Donepezil hydrochloride from isopropyl alcohol,
 (3-19) Conversion of polymorph (I), (II), (IV), (V) or amorphous form of Donepezil hydrochloride in solvent; or
 20 (3-20) According to a process (3-19) wherein the solvent is selected from methanol, ethanol, ethyl acetate or acetone.

Process (3-11) is preferred. This process is illustrated in Examples 59.

Process (3-16) is also preferred. This process is illustrated in Examples 72.

Process (3-19) is also preferred. This process is illustrated in Examples 76 to 95.

25 (4) A process for preparing the polymorph (IV) are:

- (4-2) Dissolving Donepezil in hydrochloric acid, followed by filtration of the separated crystals,
 30 (4-3) Dissolving Donepezil in hydrochloric acid, followed by addition of tetrahydrofuran,
 (4-4) Dissolving Donepezil in a mixture of water and tetrahydrofuran, followed by addition of hydrochloric acid or hydrogen chloride,
 (4-5) Dissolving Donepezil in methanol, toluene or n-hexane, followed by addition of hydrochloric acid,
 (4-6) Dissolving Donepezil in a mixture of methanol and hydrochloric acid,
 35 (4-7) Dissolving Donepezil in water, followed by addition of hydrochloric acid or hydrogen chloride,
 (4-8) Recrystallization of Donepezil hydrochloride from water,
 (4-9) Humidification of the polymorph (II) of Donepezil hydrochloride; or
 (4-10) Humidification of the amorphous form of Donepezil hydrochloride.

Process (4-4) is preferred. This process is illustrated in Example 101.

40 Process (4-8) is also preferred. This process is illustrated in Example 106.

(5) A process for preparing the polymorph (V) is:

- 45 (5-1) Heating the polymorph (IV) of Donepezil hydrochloride.

Process (5-1) is illustrated in Example 109.

50 **[0034]** In the aforementioned process (3-6) for obtaining the polymorph (III), comprising from dissolving Donepezil in ethanol, followed by addition of hydrochloric acid or hydrogen chloride, and then adding diisopropyl ether, the preferred crystallization time depends on stirring velocity, volume of solvent e.t.c. However, higher temperature shortens the crystallization time. These changes are illustrated in Example 96 to 98.

[0035] The invention provides a method for treating a disease accompanied by acetylcholinesterase activity by administering to a human patient a pharmacologically effective amount of the Donepezil hydrochloride in the form of polymorph as above for inhibiting the acetylcholinesterase activity.

55 **[0036]** The invention further provides a therapeutical composition which comprises a pharmacologically effective amount of Donepezil hydrochloride in the form of polymorph as above and a pharmacologically acceptable carrier.

[0037] The compound in the form of polymorph of the present invention is effective for treatment, prevention, remission, improvement, etc. of various kinds of senile dementia, particularly senile dementia of the Alzheimer type; cere-

brovascular diseases accompanying cerebral apoplexy, e.g. cerebral hemorrhage or cerebral infarcts, cerebral arteriosclerosis, head injury, etc.; and aprosexia, disturbance of speech, hypobulia, emotional changes, attention deficit/hyperactivity disorder, recent memory disturbance, hallucinatory-paranoid syndrome, behavioral changes, etc. accompanying encephalitis, cerebral palsy, etc.

[0038] Further, the compound in the form of polymorph of the present invention has a strong and highly selective acetylcholinesterase action, which renders the compound of the present invention useful also as a pharmaceutical based on this kind of action.

[0039] Specifically, the compound in the form of polymorph of the present invention is effective for, for example, Huntington's chorea, Pick's disease and delayed ataxia or tardive dyskinesia other than senile dementia of the Alzheimer type.

[0040] When the compound in the form of polymorph of the present invention is used as a pharmaceutical for these diseases, it may be orally or parenterally administered. In general, it is parenterally administered in the form of injections, such as intravenous, subcutaneous, and intramuscular injections, suppositories, or sublingual tablets. The dose will remarkably vary depending upon the symptom; age, sex, weight, and sensitivity of patients; method of administration; time and intervals of administration and properties, dispensing, and kind of pharmaceutical preparations; kind of effective ingredients, etc., so that there is no particular limitation with respect to the dose. Normally the compound may be administered in a dose of about 1.0 to 300 mg, preferably 1 to 100 mg, per day per adult, ordinarily in one to four portions.

[0041] Pharmaceutical preparations in the dosage form of, e.g., injections, suppositories, sublingual tablets, tablets, and capsules are prepared according to a method which is commonly accepted in the art.

[0042] In preparing injections, the effective ingredient is blended, if necessary, with a pH modifier, a buffer, a suspending agent, a solubilizing agent, a stabilizer, a tonicity agent, a preservative, etc., followed by preparation of an intravenous, subcutaneous, or intramuscular injection according to an ordinary method. In this case, if necessary, it is possible to lyophilize these preparations according to an ordinary method.

[0043] Examples of the suspending agents include methylcellulose, Polysorbate 80, hydroxyethylcellulose, acacia, powdered tragacanth, sodium carboxymethylcellulose, and polyoxyethylene sorbitan monolaurate.

[0044] Examples of the solubility agent include polyoxyethylene hydrogenated castor oil, Polysorbate 80, nicotinamide, polyoxyethylene sorbitan monolaurate, Macrogol, and an ethyl ester of castor oil fatty acid.

[0045] Examples of the stabilizer include sodium sulfite, sodium metabisulfite and ether, and examples of the preservative include methyl p-hydroxybenzoate, ethyl p-hydroxybenzoate, sorbic acid, phenol, cresol, and chlorocresol.

[Explanation of the drawings]

[0046]

Fig. 1 is powder X-ray diffraction pattern of the polymorph (I).

Fig. 2 is powder X-ray diffraction pattern of the polymorph (II).

Fig. 3 is powder X-ray diffraction pattern of the polymorph (III).

Fig. 4 is powder X-ray diffraction pattern of the polymorph (IV).

Fig. 5 is powder X-ray diffraction pattern of the Amorphous form.

Fig. 6 is infrared absorption in potassium bromide of the polymorph (I).

Fig. 7 is infrared absorption in potassium bromide of the polymorph (II).

Fig. 8 is infrared absorption in potassium bromide of the polymorph (III).

Fig. 9 is infrared absorption in potassium bromide of the polymorph (IV).

Fig. 10 is infrared absorption in potassium bromide of the amorphous form.

Fig. 11 is thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (I).

Fig. 12 is thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (II).

Fig. 13 is thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (III).

Fig. 14 is thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (IV).

Fig. 15 is thermogravimetric and differential thermal analysis (TG-DTA) of the amorphous form.

Fig. 16 is change of impurity contents for each polymorphs and amorphous form stored at -20°C.

Fig. 17 is change of impurity contents for each polymorphs and amorphous form stored at 40°C.

Fig. 18 is change of impurity contents for each polymorphs and amorphous form stored at 60°C.

Fig. 19 is change of impurity contents for each polymorphs and amorphous form stored at 80°C.

Fig. 20 is water contents for each polymorphs and amorphous form stored at 25°C under various relative humidity condition.

Fig. 21 is clear powder X-ray diffraction pattern of the polymorph (I).

Fig. 22 is clear powder X-ray diffraction pattern of the polymorph (II).

Fig. 23 is clear powder X-ray diffraction pattern of the polymorph (III).
 Fig. 24 is clear powder X-ray diffraction pattern of the polymorph (IV).
 Fig. 25 is clear powder X-ray diffraction pattern of the polymorph (V).
 Fig. 26 is clear powder X-ray diffraction pattern of the Amorphous form.
 Fig. 27 is clear infrared absorption in potassium bromide of the polymorph (I).
 Fig. 28 is clear infrared absorption in potassium bromide of the polymorph (II).
 Fig. 29 is clear infrared absorption in potassium bromide of the polymorph (III).
 Fig. 30 is clear infrared absorption in potassium bromide of the polymorph (IV).
 Fig. 31 is clear infrared absorption in potassium bromide of the polymorph (V).
 Fig. 32 is clear infrared absorption in potassium bromide of the amorphous form.
 Fig. 33 is clear thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (I).
 Fig. 34 is clear thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (II).
 Fig. 35 is clear thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (III).
 Fig. 36 is clear thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (IV).
 Fig. 37 is clear thermogravimetric and differential thermal analysis (TG-DTA) of the polymorph (V).
 Fig. 38 is clear thermogravimetric and differential thermal analysis (TG-DTA) of the amorphous form.

[0047] The products shown in Figs. 1 to 15 were again confirmed experimentally to obtain a more clear chart.

[Examples]

[0048] The present invention will now be described in more detail with reference to the following Examples. It is needless to say that the technical scope of the present invention is not limited to these Examples.

Examples 1 to 8 relates to processes for preparing a polymorph (I).
 Examples 9 to 15, processes for preparing a polymorph (II).
 Examples 16 to 27, processes for preparing a polymorph (III).
 Examples 28, process for preparing a polymorph (IV).
 Reference Examples 1, process for preparing a amorphous form of Donepezil hydrochloride.

Example 1; Polymorph (I) of Donepezil hydrochloride

[0049] One gram of Donepezil hydrochloride was dissolved in 5 ml of methanol. Adding 10 ml of diisopropyl ether and stirring the mixture in a bath containing ice water, filtration of the separated crystal and drying under atmosphere afforded 0.9 g of the title compound.

Example 2; Polymorph (I) of Donepezil hydrochloride

[0050] One gram of Donepezil hydrochloride was dissolved in 5 ml of methanol under heating. After cooling to room temperature, 10 ml of isopropyl ether was added. Stirring was continued for 30 minutes at room temperature, then filtration of the separated crystal and drying under atmosphere afforded 0.9 g of the title compound.

Example 3; Polymorph (I) of Donepezil hydrochloride

[0051] One gram of Donepezil hydrochloride was dissolved in 5 ml of methanol under heating. After initiation of cooling the solution, it started to separate crystal at 15°C of the inner temperature. After 10 minutes, 10 ml of isopropyl ether was added. Stirring was continued for 1 hour at room temperature, then filtration of the separated crystal and drying under atmosphere afforded 0.9 g of the title compound.

Example 4; Polymorph (I) of Donepezil hydrochloride

[0052] 5 g of Donepezil hydrochloride was dissolved in 25 ml of methanol under heating, followed by cooling the mixture in a bath containing ice water. Filtration of the separated crystal and drying under atmosphere afforded 4.6 g of the title compound.

Example 5; Polymorph (I) of Donepezil hydrochloride

[0053] 0.3 g of Donepezil was dissolved in 1.5 ml of methanol, followed by addition of 0.97 ml of 10%-hydrochloric

acid in methanol mixture. Filtration of the separated crystal and drying under atmosphere afforded 0.2 g of the title compound.

Example 6; Polymorph (I) of Donepezil hydrochloride

[0054] 0.3 g of Donepezil was dissolved in 3 ml of ethanol under heating, followed by addition of 3 ml of diisopropyl ether and 0.79 ml of 10%-hydrochloric acid in methanol mixture. Filtration of the separated crystal and drying under atmosphere afforded 0.2 g of the title compound.

Example 7; Polymorph (I) of Donepezil hydrochloride

[0055] 10 g of Donepezil was dissolved in 100 ml of ethanol under heating. Under stirring, a mixture of concentrated hydrochloric acid (3.1 g) and ethanol (28 ml) was added hereinto, followed by addition of 150 ml of diisopropyl ether. Filtration of the crystals after 10 seconds from the separation and drying under atmosphere afforded 9.36g of the title compound with a yield of 85.4%, a water content of 5.17% and melting point of 225-226° C (Decomposition).

Example 8; Polymorph (I) of Donepezil hydrochloride

[0056] 10 g of Donepezil hydrochloride was dissolved in 50 ml of methanol under heating. Under stirring in a bath containing ice water, 600 ml of diethyl ether was added. Stirring was continued for 1 hour in the same condition, then filtration of the crystals and drying under atmosphere afforded 10.0 g of the title compound.

Example 9; Polymorph (II) of Donepezil hydrochloride

[0057] 13.7 g of Donepezil and 4.4ml of hydrochloric acid were dissolved in 100 ml of ethanol under heating. Under stirring at room temperature, 200 ml of diisopropyl ether was added. Filtration of the crystals and drying under vacuum afforded 11.2 g of the title compound.

Example 10; Polymorph (II) of Donepezil hydrochloride

[0058] 50 g of Donepezil was dissolved in 200 ml of ethanol under heating. After cooling to room temperature, 27.3 g of 18%-hydrogen chloride in ethanol solution was added. After setting calmly for 1 hour, the mixture was concentrated under vacuum, then drying the obtained crystal under atmosphere afforded 55.0 g of the title compound.

Example 11; Polymorph (II) of Donepezil hydrochloride

[0059] 0.5 g of Donepezil was dissolved in 5 ml of ethanol under heating. Under stirring at room temperature, 1.31 ml of 10%-concentrated hydrochloric acid in ethanol was added, followed by addition of 5 ml of diisopropyl ether. Filtration of the crystals after 10 minutes from the separation and drying under atmosphere afforded 0.4 g of the title compound.

Example 12; Polymorph (II) of Donepezil hydrochloride

[0060] 5.6 g of Donepezil hydrochloride was dissolved in 30 ml of ethanol, followed by addition of 100 ml of diisopropyl ether. The mixture was cooled in a bath containing ice water. Then filtration of the crystals and drying at 50° C for three days afforded 4.9 g of the title compound.

Example 13; Polymorph (II) of Donepezil hydrochloride

[0061] 23.3 g of Donepezil hydrochloride was dissolved in 250 ml of ethanol under heating. Under stirring in a bath containing ice water, 600 ml of diethyl ether was added. After setting calmly for 3 hours, filtration of the crystals and drying at 85° C for 22 hours afforded 22.7 g of the title compound.

Example 14; Polymorph (II) of Donepezil hydrochloride

[0062] 10 g of Donepezil was dissolved in 100 ml of ethanol under heating. Under stirring, 150 ml of a mixture of concentrated hydrochloric acid (3.1 g) and ethanol (28 ml) was added, followed by addition of 150 ml of diisopropyl ether. Filtration of the crystals after 15 minutes from the separation and drying under atmosphere afforded 9.0 g of the

title compound with a yield of 82.1% and melting point of 224-226° C (Decomposition).

Example 15; Polymorph (II) of Donepezil hydrochloride

[0063] 40.0 g of Donepezil hydrochloride was dissolved in 700 ml of ethanol under heating. Under cooling in a bath containing ice water, 500 ml of diisopropyl ether was added, and crystallization was done by rubbing the flask wall with spatula.

[0064] Then filtration of the crystals and drying at 50° C for 12 hours afforded 31.4 g of the title compound.

Example 16; Polymorph (III) of Donepezil hydrochloride

[0065] 161 g of Donepezil hydrochloride was dissolved in 2,000 ml of ethanol under heating. After cooling to room temperature, 5,000 ml of diethyl ether was added under stirring. Then filtration of the crystals and drying at 35° C for 12 hours afforded 120 g of the title compound with a yield of 74.5% and a water content of 0.15%.

Example 17; Polymorph (III) of Donepezil hydrochloride

[0066] 308 g of Donepezil was dissolved in 700 ml of ethanol. Under stirring, 230 ml of 10%-hydrogen chloride in ethanol solution and 5000 ml of diethyl ether were added successively. Filtration of the crystals and drying at 50° C for 1 hour, then at 60° C for 30 minutes, then at 85° C for 12 hours afforded 269g of the title compound.

Example 18; Polymorph (III) of Donepezil hydrochloride

[0067] 59 g of Donepezil was dissolved in 590 ml of ethanol. Under cooling in a bath containing ice water, 17.8 g of concentrated hydrochloric acid and 885 ml of diisopropyl ether were added successively. After stirring over night at room temperature, filtration of the crystals and drying at 55° C for 22 hours afforded 62 g of the title compound.

Example 19; Polymorph (III) of Donepezil hydrochloride

[0068] 5 g of Donepezil hydrochloride was dissolved in 100 ml of ethanol under heating. After cooling to room temperature, 100 ml of n-hexane was added hereinto under stirring, followed by cooling in a bath containing ice water. Stirring was continued for 1 hour. Filtration of the crystals and drying at room temperature afforded 4 g of the title compound.

Example 20; Polymorph (III) of Donepezil hydrochloride

[0069] One gram of Donepezil hydrochloride was dissolved in 15 ml of dichloromethane under heating. After cooling to room temperature, 15 ml of n-hexane was added hereinto under stirring, followed by cooling in a bath containing ice water. Stirring was continued for 1 hour. Filtration of the crystals and drying at room temperature afforded 0.9 g of the title compound.

Example 21; Polymorph (III) of Donepezil hydrochloride

[0070] 0.5 g of Donepezil was dissolved in 10 ml of acetone under heating. Under stirring at room temperature, 0.13 ml of concentrated hydrochloric acid was added hereinto. Stirring was continued for 30 minutes. Filtration of the crystals and drying at 85°C for 16 hours afforded 0.5 g of the title compound.

Example 22; Polymorph (III) of Donepezil hydrochloride

[0071] 0.3 g of Donepezil was dissolved in 3 ml of ethyl acetate under heating. Under stirring at room temperature, 0.79 ml of 10%-hydrogen chloride in ethanol solution was added. Filtration of the crystals and drying at 85° C for 3 hours, then at 70° C for 16 hours afforded 0.3g of the title compound.

Example 23; Polymorph (III) of Donepezil hydrochloride

[0072] 10 g of Donepezil was dissolved in 100 ml of ethanol under heating. Under stirring, a mixture of 3.1 g of concentrated hydrochloric acid and 28 ml of ethanol and then 150 ml of diisopropyl ether were added successively. Stirring was continued for 1 hour from the separation of crystals. Filtration of the crystals and drying at room temperature

afforded 9.86g of the title compound with a yield of 90%, a water content of 0.26% and melting point of 229-231° C (Decomposition).

Example 24; Polymorph (III) of Donepezil hydrochloride

[0073] 5.0 g of The polymorph (I) of Donepezil hydrochloride was spread on a laboratory dish (Schale) and was allowed to stand for 7 days under ventilation of air at 85° C. 4.9 g of the title compound was obtained.

Example 25; Polymorph (III) of Donepezil hydrochloride

[0074] 5.0g of the polymorph (I) of Donepezil hydrochloride was spread on a laboratory dish (Schale) and was allowed to stand for 2 days under ventilation of air at 85° C, then for 3 days under ventilation of air at 105°C. 4.8 g of the title compound was obtained.

Example 26; Polymorph (III) of Donepezil hydrochloride

[0075] 5.0 g of The polymorph (I) of Donepezil hydrochloride was spread on a laboratory dish (Schale) and was allowed to stand for 5 days under ventilation of air at 85° C. 4.9 g of the title compound was obtained.

Example 27; Polymorph (III) of Donepezil hydrochloride

[0076] 5.0 g of the polymorph (I) of Donepezil hydrochloride was spread on a laboratory dish (Schale) and was allowed to stand for 3 days under ventilation of air at 105° C. 4.9 g of the title compound was obtained.

Example 28; Polymorph (IV) of Donepezil hydrochloride

[0077] 15.0 g of the polymorph (II) of Donepezil hydrochloride was spread on a laboratory dish (Schale) and was continued to stand for 2 weeks under atmosphere having a relative humidity of 100%. 14.8 g of the title compound was obtained with melting point of 226-228°C (Decomposition).

Reference Examples 1; Amorphous form of Donepezil hydrochloride

[0078] 15.0 g of Donepezil hydrochloride was dissolved in 300 ml of water. The solution was frozen in a bath containing dry ice and acetone and freezing-dried (lyophilized) for 4 days at -82°C. 14.8 g of the title compound was obtained.

[0079] The present invention will be futhermore described in detail with reference to the following Examples.

Examples 29 to 44, processes for preparing a polymorph (I).

Examples 45 to 57, processes for preparing a polymorph (II).

Examples 58 to 98, process for preparing a polymorph (III).

Examples 99 to 108 relates to processes for preparing a polymorph (IV).

Example 109, process for preparing a polymorph (V).

Example 29; Polymorph (I) of Donepezil hydrochloride

[0080] 1.0 g of Donepezil was dissolved in 4 ml of methanol under heating at 40°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of methanol was added hereinto at 10°C inner temperature. It continued stirring for 90 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.43 g of the title compound. (water content: 5.33%)

Example 30; Polymorph (I) of Donepezil hydrochloride

[0081] 1.0 g of Donepezil was dissolved in 4 ml of methanol under heating at 40°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of methanol was added hereinto. After 5 minutes, 30 ml of tert-butyl methyl ether (hereinafter, abbreviated as TBME) was added at 3°C inner temperature. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.10 g of the title compound. (water content: 5.60%)

Example 31; Polymorph (I) of Donepezil hydrochloride

[0082] 1.0 g of Donepezil was dissolved in 4 ml of methanol under heating at 40°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of methanol was added hereinto. After 5 minutes, 30 ml of diisopropyl ether (hereinafter, abbreviated as IPE) was added at 3°C inner temperature. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.13 g of the title compound. (water content: 5.50%)

Example 32; Polymorph (I) of Donepezil hydrochloride

[0083] 1.0 g of Donepezil was dissolved in 4 ml of methanol under heating at 40°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of methanol was added hereinto at 12°C inner temperature. After 7 minutes, 30 ml of ethyl acetate was added at 3°C inner temperature successively. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.71 g of the title compound. (water content: 5.22%)

Example 33; Polymorph (I) of Donepezil hydrochloride

[0084] 1.0 g of Donepezil was dissolved in 4 ml of ethanol under heating at 40°C. The solution was cooled in an iced water bath. After cooling for 5 minutes, 0.31 g of concentrated hydrochloric acid in 1 ml of ethanol was added hereinto. It continued stirring for 30 minutes in an iced water bath. A small portion of the polymorph (I) of Donepezil hydrochloride was added hereinto. It continued stirring for further 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.70 g of the title compound. (water content: 5.33%)

Example 34; Polymorph (I) of Donepezil hydrochloride

[0085] 1.0 g of Donepezil was dissolved in 4 ml of tetrahydrofuran (hereinafter, abbreviated as THF) at 24°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of THF was added hereinto. It continued stirring for 40 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.00 g of the title compound. (water content: 5.67%)

Example 35; Polymorph (I) of Donepezil hydrochloride

[0086] 1.0 g of Donepezil was dissolved in 9 ml of acetonitrile under heating at 40°C. The solution was cooled in an iced water bath. After cooling for 2 minutes, 0.31 g of concentrated hydrochloric acid in 1 ml of acetonitrile was added hereinto. It continued stirring for 50 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.63 g of the title compound. (water content: 5.59%)

Example 36; Polymorph (I) of Donepezil hydrochloride

[0087] 4.0 g of Donepezil was dissolved in 20 ml of methanol under heating at 40°C. The solution was cooled in an iced water bath. Hydrogen chloride gas was blown hereinto at 3°C inner temperature, until the atmosphere turns acidic. It continued stirring for 20 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 3.40 g of the title compound. (water content: 5.19%)

Example 37; Polymorph (I) of Donepezil hydrochloride

[0088] 10.0 g of Donepezil hydrochloride was dissolved in 60 ml of methanol under reflux. Heating was ceased. 120 ml of IPE was added hereinto at 60°C inner temperature. It continued stirring for 20 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 9.80 g of the title compound. (water content: 5.87%)

Example 38; Polymorph (I) of Donepezil hydrochloride

[0089] 3.0 g of Donepezil hydrochloride was dissolved in 18 ml of methanol under reflux. Heating was ceased. 36 ml of IPE was added hereinto at 54°C inner temperature. It continued stirring for 20 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 2.95 g of the title compound. (water content: 5.55%)

[0090] Elementary Analysis:

EP 1 211 243 A1

| | C | H | N | Cl |
|----------|-------|------|------|------|
| found(%) | 65.55 | 7.53 | 3.05 | 8.16 |

Example 39; Polymorph (I) of Donepezil hydrochloride

[0091] 1.0 g of Donepezil hydrochloride was dissolved in 5 ml of methanol under reflux. The solution was cooled in an iced water bath. 30 ml of TBME was added hereinto at 3°C inner temperature. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.00 g of the title compound. (water content: 5.40%)

Example 40; Polymorph (I) of Donepezil hydrochloride

[0092] 1.0 g of Donepezil hydrochloride was dissolved in 5 ml of methanol under reflux. The solution was cooled in an iced water bath. After cooling for 2 minutes, it started to separate crystals. 1 Minutes passed from the initiation of separation, 30 ml of ethyl acetate was added hereinto. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.00 g of the title compound. (water content: 5.60%)

Example 41; Polymorph (I) of Donepezil hydrochloride

[0093] 1.0 g of Donepezil hydrochloride was dissolved in 5 ml of methanol under reflux. The solution was cooled in an iced water bath. 30 ml of n-hexane was added hereinto at 3°C inner temperature. It continued stirring for 70 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.89 g of the title compound. (water content: 5.66%)

Example 42; Polymorph (I) of Donepezil hydrochloride

[0094] 1.0 g of Donepezil hydrochloride was dissolved in 20 ml of ethanol under reflux. The solution was cooled in an iced water bath. It continued stirring for 70 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.48 g of the title compound. (water content: 5.72%)

Example 43; Polymorph (I) of Donepezil hydrochloride

[0095] 1.0 g of Donepezil hydrochloride was dissolved in 25 ml of ethanol under reflux. The solution was cooled in an iced water bath. 50 ml of IPE was added at 3°C inner temperature. It continued stirring for 5 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.86 g of the title compound. (water content: 5.32%)

Example 44; Polymorph (I) of Donepezil hydrochloride

[0096] 1.0 g of Donepezil hydrochloride was dissolved in 20 ml of ethanol under reflux. The solution was cooled in an iced water bath. 30 ml of TBME was added hereinto at 3°C inner temperature. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.00 g of the title compound. (water content: 5.33%)

Example 45; Polymorph (II) of Donepezil hydrochloride

[0097] 1.0 g of Donepezil was dissolved in 4 ml of ethanol under heating at 40°C. The solution was cooled in an iced water bath. After cooling for 4 minutes, 0.31 g of concentrated hydrochloric acid in 1 ml of ethanol was added hereinto. After 3 minutes, 30 ml of TBME was added hereinto. It continued stirring for 50 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.08 g of the title compound. (water content: 1.78%)

Example 46; Polymorph (II) of Donepezil hydrochloride

[0098] 1.0 g of Donepezil was dissolved in 9 ml of isopropyl alcohol (hereinafter, abbreviated as IPA) under heating at 40°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of IPA was added hereinto. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed

EP 1 211 243 A1

by drying afforded 0.87 g of the title compound. (water content: 1.10%)

Example 47; Polymorph (II) of Donepezil hydrochloride

[0099] 1.0 g of Donepezil was dissolved in 9 ml of acetone at 19°C. 0.31 g of Concentrated hydrochloric acid in 1 ml of acetone was added hereinto. It continued stirring for 5 minutes at room temperature. Filtration of the separated crystals followed by drying afforded 0.87 g of the title compound. (water content: 0.83%)

Example 48; Polymorph (II) of Donepezil hydrochloride

[0100] 1.0 g of Donepezil was dissolved in 9 ml of acetone at 24°C. 0.31 g of Concentrated hydrochloric acid in 1 ml of acetone was added hereinto. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.92 g of the title compound. (water content: 0.61%)

Example 49; Polymorph (II) of Donepezil hydrochloride

[0101] 1.0 g of Donepezil was dissolved in 9 ml of THF at 24°C. 0.31 g of Concentrated hydrochloric acid in 1 ml of THF was added hereinto. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.09 g of the title compound. (water content: 0.78%)

Example 50; Polymorph (II) of Donepezil hydrochloride

[0102] 3.0 g of Donepezil was dissolved in 30 ml of acetone at 21°C. Hydrogen chloride gas was blown hereinto at room temperature, until the atmosphere turns acidic. After stirring for 3 minutes, filtration of the separated crystals followed by drying afforded 2.80 g of the title compound. (water content: 2.78%)

Example 51; Polymorph (II) of Donepezil hydrochloride

[0103] 4.0 g of Donepezil was dissolved in 20 ml of methylene chloride at 18°C. The solution was cooled in an iced water bath. Hydrogen chloride gas was blown hereinto at 4°C inner temperature, until the atmosphere turns acidic. Argon gas was blown hereinto. After stirring for 2 hours in an iced water bath, filtration of the separated crystals followed by drying afforded 4.09 g of the title compound. (water content: 0.81%)

Example 52; Polymorph (II) of Donepezil hydrochloride

[0104] 1.0 g of Donepezil hydrochloride was dissolved in 20 ml of ethanol under reflux. The solution was cooled in an iced water bath. 30 ml of TBME was added hereinto at 20°C inner temperature. It continued stirring for 3 hours in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.92 g of the title compound. (water content: 0.79%)

Example 53; Polymorph (II) of Donepezil hydrochloride

[0105] 1.0 g of Donepezil hydrochloride was dissolved in 20 ml of ethanol under reflux. The solution was cooled to room temperature in an iced water bath. 30 ml of TBME was added hereinto at 20°C inner temperature. It continued stirring for 20 minutes at room temperature. Filtration of the separated crystals followed by drying afforded 0.80 g of the title compound. (water content: 0.52%)

Example 54; Polymorph (II) of Donepezil hydrochloride

[0106] 10.0 g of Donepezil hydrochloride was dissolved in 100 ml of ethanol under reflux. Under stirring, the solution was added into 200 ml of IPE cooled in an iced water bath. It continued stirring for 5 minutes Filtration of the separated crystals followed by drying afforded 9.40 g of the title compound. (water content: 0.19%)

[0107] Elementary Analysis:

| | C | H | N | Cl |
|----------|-------|------|------|------|
| found(%) | 69.12 | 7.20 | 3.32 | 8.61 |

EP 1 211 243 A1

Example 55; Polymorph (II) of Donepezil hydrochloride

[0108] 1.0 g of Donepezil hydrochloride was dissolved in 20 ml of methylene chloride at 18°C. The solution was cooled in an iced water bath. 30 ml of TBME was added hereinto. After stirring for 3 minutes in an iced water bath, filtration of the separated crystals followed by drying afforded 0.88 g of the title compound. (water content: 2.33%)

Example 56; Polymorph (II) of Donepezil hydrochloride

[0109] 2.0 g of The polymorph (I) of Donepezil hydrochloride was spread on an laboratory dish (Schale) and was allowed to stand for 16 hours under reduced pressure at 80°C. 1.89 g of the title compound was obtained. (water content: 0.22%)

Example 57; Polymorph (II) of Donepezil hydrochloride

[0110] 2.0 g of The amorphous form of Donepezil hydrochloride was spread on an laboratory dish (Schale) and was allowed to stand for 16 hours under reduced pressure at 80°C. 1.98 g of the title compound was obtained. (water content: 1.15%)

Example 58; Polymorph (III) of Donepezil hydrochloride

[0111] 1.0 g of Donepezil was dissolved in 4 ml of methanol under heating at 40°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of methanol was added hereinto at 7°C inner temperature. After 5 minutes stirring, 30 ml of acetone was added hereinto successively. It continued stirring for 1 hour in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.44 g of the title compound. (water content: 0.11%)

Example 59; Polymorph (III) of Donepezil hydrochloride

[0112] 1.0 g of Donepezil was dissolved in 4 ml of ethanol under heating at 40°C. The solution was cooled to room temperature in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of ethanol was added hereinto. After 5 minutes, 30 ml of TBME was added hereinto at 22°C inner temperature. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded 1.09 g of the title compound. (water content: 0.19%)

[0113] Elementary Analysis:

| | C | H | N | Cl |
|----------|-------|------|------|------|
| found(%) | 69.27 | 7.23 | 3.34 | 8.58 |

Example 60; Polymorph (III) of Donepezil hydrochloride

[0114] 1.0 g of Donepezil was dissolved in 9 ml of ethyl acetate under heating at 40°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of ethyl acetate was added hereinto. It continued stirring for 30 minutes at room temperature. Filtration of the separated crystals followed by drying afforded 1.08 g of the title compound. (water content: 0.21%)

Example 61; Polymorph (III) of Donepezil hydrochloride

[0115] 1.0 g of Donepezil was almost dissolved in 9 ml of acetonitrile at 21°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of acetonitrile was added hereinto. The solution became homogeneous. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded 1.05 g of the title compound. (water content: 0.05%)

Example 62; Polymorph (III) of Donepezil hydrochloride

[0116] 1.0 g of Donepezil was dissolved in 9 ml of acetone under heating at 40°C. The solution was cooled to room temperature in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of acetone was added hereinto. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded 1.08

g of the title compound. (water content: 0.03%)

Example 63; Polymorph (III) of Donepezil hydrochloride

[0117] 1.0 g of Donepezil was dissolved in the mixture comprising from 29 ml of acetone and 1 ml of ion exchange purified water under stirring at 20°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of acetone was added hereinto. After stirring for 30 minutes in an iced water bath, filtration of the separated crystals followed by drying afforded 0.83 g of the title compound. (water content: 0.56%)

Example 64; Polymorph (III) of Donepezil hydrochloride

[0118] 1.0 g of Donepezil was dissolved in 9 ml of THF at 24°C. 0.31 g of Concentrated hydrochloric acid in 1 ml of THF was added hereinto. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.09 g of the title compound. (water content: 0.54%)

Example 65; Polymorph (III) of Donepezil hydrochloride

[0119] 1.0 g of Donepezil was dissolved in 4 ml of DMF under heating at 40°C. The solution was cooled in an iced water bath. 0.31 g of Concentrated hydrochloric acid in 1 ml of DMF was added hereinto. It continued stirring for 30 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded 1.08 g of the title compound. (water content: 0.72%)

Example 66; Polymorph (III) of Donepezil hydrochloride

[0120] 1.0 g of Donepezil was almost dissolved in 9 ml of DMSO at 20°C. 0.31 g of Concentrated hydrochloric acid in 1 ml of DMSO was added hereinto. After stirring for 1 hour at room temperature, 30 ml of TBME was added hereinto. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded 0.97 g of the title compound. (water content: 0.25%)

Example 67; Polymorph (III) of Donepezil hydrochloride

[0121] 2.0 g of Donepezil was dissolved in 20 ml of ethanol under heating at 40°C. The solution was cooled to room temperature in an iced water bath. Hydrogen chloride gas was blown hereinto at 24°C inner temperature, until the atmosphere turns acidic. After stirring for 10 minutes, 50 ml of TBME was added. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded 2.11 g of the title compound. (water content: 0.07%)

Example 68; Polymorph (III) of Donepezil hydrochloride

[0122] 2.0 g of Donepezil was dissolved in 30 ml of toluene at 20°C. Hydrogen chloride gas was blown hereinto at room temperature, until the atmosphere turns acidic. Argon gas was blown hereinto. After stirring for 30 minutes, filtration of the separated crystals followed by drying afforded 2.21 g of the title compound. (water content: 0.65%)

Example 69; Polymorph (III) of Donepezil hydrochloride

[0123] 1.0 g of Donepezil hydrochloride was dissolved in 5 ml of methanol under reflux. Heating was ceased. After stirring over night at room temperature, it continued stirring 2 hours additionally in an iced water. Filtration of the separated crystals followed by drying afforded 0.90 g of the title compound. (water content: 0.05%)

Example 70; Polymorph (III) of Donepezil hydrochloride

[0124] 1.0 g of Donepezil hydrochloride was dissolved in 5 ml of methanol under reflux. The solution was cooled, then 30 ml of TBME was added hereinto at 40°C inner temperature. It continued stirring for 3 hours at 40°C. The mixture was cooled in an iced water bath, and was kept for 45 minutes under stirring. Filtration of the separated crystals followed by drying afforded 0.94 g of the title compound. (water content: 0.13%)

EP 1 211 243 A1

Example 71; Polymorph (III) of Donepezil hydrochloride

[0125] 1.0 g of Donepezil hydrochloride was dissolved in 5 ml of methanol under reflux. The solution was cooled in an iced water bath. After 4 minutes, it started to separate crystals. 30 ml of Acetonitrile was added hereinto, then the crystal was dissolved immediately. After stirring over night at room temperature, filtration of the separated crystals followed by drying afforded 0.10 g of the title compound. (water content: 0.09%)

Example 72; Polymorph (III) of Donepezil hydrochloride

[0126] 1.0 g of Donepezil hydrochloride was dissolved in 20 ml of ethanol under reflux. The solution was cooled, then 30 ml of TBME was added hereinto at 40°C inner temperature. It continued stirring for 3 hours at 40°C. The mixture was cooled in an iced water bath, and was kept for 20 minutes under stirring. Filtration of the separated crystals followed by drying afforded 0.97 g of the title compound. (water content: 0.14%)

Example 73; Polymorph (III) of Donepezil hydrochloride

[0127] 1.0 g of Donepezil hydrochloride was dissolved in 120 ml of IPA under reflux. Heating was ceased. It was kept under stirring over night at room temperature. It continued stirring for 1 hour in an iced water bath. Filtration of the separated crystals followed by drying afforded 0.92 g of the title compound. (water content: 0.21%)

Example 74; Polymorph (III) of Donepezil hydrochloride

[0128] 1.0 g of Donepezil hydrochloride was dissolved in 15 ml of N,N-dimethylformamide at 100°C. Heating was ceased, and the solution was kept over night under stirring. 30 ml of TBME was added hereinto at 20°C inner temperature. The mixture was cooled in an iced water bath, and was kept for 3 hours under stirring. Filtration of the separated crystals followed by drying afforded 0.90 g of the title compound. (water content: 0.10%)

Example 75; Polymorph (III) of Donepezil hydrochloride

[0129] 1.0 g of Donepezil hydrochloride was dissolved in 15 ml of dimethylsulfoxide (hereinafter, abbreviated as DMSO) under heating at 80°C. The solution was cooled to room temperature in an iced water bath. 30 ml of TBME was added hereinto at 20°C inner temperature. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded 1.01 g of the title compound. (water content: 0.08%)

Example 76 to 95; Polymorph (III) of Donepezil hydrochloride

[0130] 1.0 g of the material shown in the below table was suspended in 10 ml of the solvent shown in the below table at room temperature. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded the title compound.

| Example | Material | Solvent | Yield |
|---------|----------------|---------------|-------|
| 76 | polymorph (I) | methanol | 0.91g |
| 77 | polymorph (I) | ethanol | 0.94g |
| 78 | polymorph (I) | ethyl acetate | 0.93g |
| 79 | polymorph (I) | acetone | 0.93g |
| 80 | polymorph (II) | methanol | 0.93g |
| 81 | polymorph (II) | ethanol | 0.95g |
| 82 | polymorph (II) | ethyl acetate | 0.95g |
| 83 | polymorph (II) | acetone | 0.94g |
| 84 | polymorph (IV) | methanol | 0.92g |
| 85 | polymorph (IV) | ethanol | 0.90g |
| 86 | polymorph (IV) | ethyl acetate | 0.92g |
| 87 | polymorph (IV) | acetone | 0.94g |
| 88 | polymorph (V) | methanol | 0.96g |
| 89 | polymorph (V) | ethanol | 0.93g |

(continued)

| Example | Material | Solvent | Yield |
|---------|----------------|---------------|-------|
| 90 | polymorph (V) | ethyl acetate | 0.97g |
| 91 | polymorph (V) | acetone | 0.92g |
| 92 | amorphous form | methanol | 0.94g |
| 93 | amorphous form | ethanol | 0.94g |
| 94 | amorphous form | ethyl acetate | 0.92g |
| 95 | amorphous form | acetone | 0.96g |

Example 96; Polymorph (III) of Donepezil hydrochloride

[0131] 10.0 g of Donepezil was dissolved in 100 ml of ethanol under heating at 40°C. The solution was cooled to room temperature. 3.01 g of Concentrated hydrochloric acid was added hereinto at 20°C inner temperature. The solution was set in an iced water bath. After 9 minutes stirring, 150 ml of isopropyl ether was added hereinto at 3°C inner temperature. It continued stirring for 300 minutes in an iced water bath. Filtration of the separated crystals followed by drying afforded the title compound.

Example 97; Polymorph (III) of Donepezil hydrochloride

[0132] 10.0 g of Donepezil was dissolved in 100 ml of ethanol under heating at 40°C. The solution was cooled to room temperature. 3.01 g of Concentrated hydrochloric acid was added hereinto at 20°C inner temperature. After 9 minutes stirring, 150 ml of isopropyl ether was added hereinto at 20°C inner temperature. It continued stirring for 120 minutes at room temperature. Filtration of the separated crystals followed by drying afforded the title compound.

Example 98; Polymorph (III) of Donepezil hydrochloride

[0133] 10.0 g of Donepezil was dissolved in 100 ml of ethanol under heating at 40°C. The solution was cooled to room temperature. 3.01 g of Concentrated hydrochloric acid was added hereinto at 20°C inner temperature. The solution was set in an oil bath, and was heated to 40°C inner temperature. 150 ml of isopropyl ether was added hereinto at 40°C inner temperature. It continued stirring for 20 minutes at 60°C inner temperature. Filtration of the separated crystals followed by drying afforded the title compound.

Example 99; Polymorph (IV) of Donepezil hydrochloride

[0134] 2.0 g of Donepezil was dissolved in the mixture comprising from 0.65 g of concentrated hydrochloric acid and 10 ml of ion exchange purified water under stirring at 22°C. After stirring at room temperature for 1 hour, filtration of the separated crystals followed by drying afforded 1.80 g of the title compound. (water content: 11.00%)

Example 100; Polymorph (IV) of Donepezil hydrochloride

[0135] 1.0 g of Donepezil was dissolved in the mixture comprising from 0.31 g of concentrated hydrochloric acid and 4 ml of ion exchange purified water under stirring at 22°C. 100 ml of THF was added hereinto. It continued stirring for 30 minutes at room temperature. Filtration of the separated crystals followed by drying afforded 1.06 g of the title compound. (water content: 11.14%)

Example 101; Polymorph (IV) of Donepezil hydrochloride

[0136] 3.0 g of Donepezil was dissolved in the mixture comprising from 88 ml of THF and 3 ml of ion exchange purified water under stirring at 20°C. 0.93 g of Concentrated hydrochloric acid in 2 ml of THF was added hereinto. After stirring for 30 minutes in an iced water bath, filtration of the separated crystals followed by drying afforded 3.21 g of the title compound. (water content: 11.34%)

[0137] Elementary Analysis:

| | C | H | N | Cl |
|----------|-------|------|------|------|
| found(%) | 61.30 | 7.76 | 2.86 | 7.68 |

Example 102; Polymorph (IV) of Donepezil hydrochloride

[0138] 1.0 g of Donepezil was dissolved in 9 ml of toluene at 22°C. 0.31 g of Concentrated hydrochloric acid in 1 ml of toluene was added hereinto. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded 1.23 g of the title compound. (water content: 11.40%)

Example 103; Polymorph (IV) of Donepezil hydrochloride

[0139] 1.0 g of Donepezil was dissolved in 9 ml of n-hexane at 21°C. 0.31 g of Concentrated hydrochloric acid in 1 ml of n-hexane was added hereinto. It continued stirring over night at room temperature. Filtration of the separated crystals followed by drying afforded 1.23 g of the title compound. (water content: 11.24%)

Example 104; Polymorph (IV) of Donepezil hydrochloride

[0140] 1.0 g of Donepezil was dissolved in the mixture comprising from 1 ml of methanol, 3 ml of ion exchange purified water and 0.93g of concentrated hydrochloric acid at 20°C. It continued stirring for 3 days at room temperature. Filtration of the separated crystals followed by drying afforded 0.83 g of the title compound. (water content: 11.04%)

Example 105; Polymorph (IV) of Donepezil hydrochloride

[0141] 2.0 g of Donepezil was suspended in 10 ml of ion exchange purified water. Hydrogen chloride gas was blown hereinto at 23°C inner temperature, until it became homogeneous. After stirring for 2.5 hours at room temperature, filtration of the separated crystals followed by drying afforded 1.77 g of the title compound. (water content: 11.10%)

Example 106; Polymorph (IV) of Donepezil hydrochloride

[0142] 1.0 g of Donepezil hydrochloride was dissolved in 5 ml of ion exchange purified water under heating at 60°C. The solution was cooled to room temperature. After stirring over night at room temperature, filtration of the separated crystals followed by drying afforded 0.92 g of the title compound. (water content: 11.12%)

Example 107; Polymorph (IV) of Donepezil hydrochloride

[0143] 1.0 g of The polymorph (II) of Donepezil hydrochloride was spread on an laboratory dish (Schale) and was allowed to stand for 24 hours under atmosphere having an relative humidity of 90%. 1.15 g of the Polymorph (IV) was obtained. (water content: 12.33%)

Example 108; Amorphous Form & Polymorph (IV) of Donepezil hydrochloride

[0144] 10.0 g of the polymorph (III) of Donepezil hydrochloride was spread on an laboratory dish (Schale, $\phi=250\text{mm}$), and was dissolved in 300 ml of ion exchange purified water at 21°C. This solution was moved into freeze-drying apparatus, and dried for 3 days to afford 9.90 g of amorphous form. This material was kept standing for 24 hours under atmosphere having an relative humidity of 90%. 11.20 g of the Polymorph (IV) was obtained. (water content: 11.21%)

Example 109; Polymorph (V) of Donepezil hydrochloride

[0145] 2.0 g of the polymorph (IV) of Donepezil hydrochloride was spread on an laboratory dish (Schale, $\phi=250\text{mm}$), and was allowed to stand for 16 hours under reduced pressure at 80°C. 1.75 g of The title compound was obtained. (water content: 0.28%)

[0146] Finally, the efficacy of the present invention in view of the stability or hygroscopicity will now be described in comparison with amorphous form of Donepezil hydrochloride. The invention provides advantageous results as follows:

(1) Stability Assay

(Method for measurement)

[0147] 10mg of each of the polymorphs (I) to (IV) of Donepezil hydrochloride was taken as a couple of samples into tubes, respectively. They were stored under the following conditions and impurity's contents were measured periodically.

EP 1 211 243 A1

| Condition | Storage Period | | | |
|-----------|----------------|---------|---------|----------|
| 80° C | 1 week | 2 weeks | | |
| 60° C | | 2 weeks | 1 month | |
| 40° C | | | 1 month | 3 months |
| -20° C | 1 week | 2 weeks | 1 month | 3 months |

(Method and condition for measurement of HPLC purity)

[0148] 10ml of the following mobile phase for HPLC was added into each tube of the aforementioned samples. Then impurity's contents were measured for each sample under the following conditions. The average was calculated from two results.

| | |
|----------------------|---|
| Column (Solid phase) | Inertsil ODS-II (4.6 mm I.D. × 150 mm) |
| Mobile Phase | CH ₃ CN/water/70% HClO ₄ (V/V/V=300:700:1) |
| Detector | UV271 nm |
| Flow rate | 1.0 ml/min. |
| Injection Volume | 5 ml |
| Column Temperature | room temperature |

Results:

[0149]

| 1) Storage at -20°C. (refer to Fig. 16) | | | | | |
|---|------|--------|---------|---------|------|
| Impurity(%) months | 0 | 1 week | 2 weeks | 1 month | 3 |
| Polymorph (I) | 0.11 | 0.12 | 0.12 | 0.13 | 0.11 |
| Polymorph (II) | 0.09 | 0.09 | 0.13 | 0.10 | 0.09 |
| Polymorph (III) | 0.15 | 0.14 | 0.13 | 0.13 | 0.15 |
| Polymorph (IV) | 0.08 | 0.08 | 0.08 | 0.08 | 0.08 |
| Amorphous | 0.12 | 0.14 | 0.15 | 0.15 | 0.14 |

| 2) Storage at 40°C. (refer to Fig. 17) | | | |
|--|------|---------|----------|
| Impurity(%) | 0 | 1 month | 3 months |
| Polymorph (I) | 0.11 | 0.12 | 0.13 |
| Polymorph (II) | 0.09 | 0.08 | 0.08 |
| Polymorph (III) | 0.15 | 0.14 | 0.15 |
| Polymorph (IV) | 0.08 | 0.08 | 0.07 |
| Amorphous | 0.12 | 0.20 | 0.45 |

| 3) Storage at 60°C. (refer to Fig. 18) | | | |
|--|------|---------|---------|
| Impurity(%) | 0 | 2 weeks | 1 month |
| Polymorph (I) | 0.11 | 0.12 | 0.13 |
| Polymorph (II) | 0.09 | 0.12 | 0.09 |
| Polymorph (III) | 0.15 | 0.14 | 0.14 |
| Polymorph (IV) | 0.08 | 0.09 | 0.13 |
| Amorphous | 0.12 | 0.30 | 0.39 |

| 4) Storage at 80°C. (refer to Fig. 19) | | | |
|--|------|--------|---------|
| Impurity(%) | 0 | 1 week | 2 weeks |
| Polymorph (I) | 0.11 | 0.12 | 0.19 |
| Polymorph (II) | 0.09 | 0.20 | 0.22 |
| Polymorph (III) | 0.15 | 0.14 | 0.14 |
| Polymorph (IV) | 0.08 | 0.09 | 0.19 |
| Amorphous | 0.12 | 2.02 | 3.29 |

[0150] It is evident from the above results that the polymorphs (I) to (IV) is superior in stability against heat to the amorphous form.

(2) Hygroscopicity Assay

(Method for measurement)

[0151] Polymorphs (I) to (IV) were stored under atmosphere having the following relative humidities at 25°C. Water contents were measured according to the general method (Karl Fischer Method) introduced by the Japanese Pharmacopoeia.

| Results (refer to Fig. 20) | | | | | |
|----------------------------|-------|-------------------|-------|-------|-----------|
| Relative Humidity | | Water contents(%) | | | |
| | | Polymorph | | | |
| (%) | (I) | (II) | (III) | (IV) | Amorphous |
| Initiation | 4.34 | 0.26 | 0.11 | 12.87 | 2.03 |
| 10.6 | 4.54 | 0.28 | 0.15 | 10.70 | 4.09 |
| 22.2 | 4.75 | 0.29 | 0.14 | 10.60 | 4.78 |
| 31.0 | 5.07 | 0.32 | 0.26 | 10.77 | 5.61 |
| 42.8 | 5.25 | 0.39 | 0.13 | 11.03 | 7.80 |
| 51.0 | 5.38 | 0.43 | 0.15 | 11.28 | 9.29 |
| 61.8 | 5.49 | 0.40 | 0.18 | 11.40 | 12.01 |
| 75.0 | 5.65 | 1.73 | 0.15 | 11.62 | 13.89 |
| 84.0 | 5.64 | 13.70 | 0.16 | 11.72 | 13.74 |
| 93.0 | 5.76 | 13.99 | 0.15 | 11.84 | 14.51 |
| 96.6 | 5.88 | 14.18 | 0.17 | 11.80 | 14.53 |
| 100.0 | 10.37 | 15.93 | 9.71 | 12.26 | 15.44 |

[0152] In the above results, the polymorphs (I) to (IV) did not show hygroscopicity until a relative humidity of 96.6%, until 75.0%, until 96.6%, until 100%, respectively. The amorphous Donepezil hydrochloride showed hygroscopicity at and thereafter 10.6%. Those experimental results show that the polymorphs of Donepezil hydrochloride (I) to (IV) have an excellent, heat stability and a low hygroscopicity.

Claims

1. Donepezil hydrochloride, 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride, in the form of polymorphs selected from the group consisting of (II), and (IV), each Polymorphs being specified by peaks at below shown diffraction degrees with the below shown intensity in terms of I/I_0 in powder X-ray diffraction pattern and the below shown absorption peaks in infrared absorption spectra in potassium bromide in terms of reciprocal centimeters:

(2) Polymorphs (II)

Peaks in the powder X-ray diffraction pattern are:

EP 1 211 243 A1

| | Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|----|---|--------------------------|
| 5 | 7.40 | 8 |
| | 9.88 | 100 |
| | 12.36 | 13 |
| | 15.54 | 40 |
| | 16.10 | 38 |
| 10 | 16.22 | 38 |
| | 16.48 | 35 |
| | 17.30 | 17 |
| | 18.04 | 20 |
| 15 | 18.44 | 17 |
| | 18.84 | 19 |
| | 19.34 | 19 |
| | 19.84 | 47 |
| | 21.16 | 24 |
| 20 | 22.40 | 19 |
| | 23.18 | 33 |
| | 24.02 | 22 |
| | 24.92 | 25 |
| 25 | 25.72 | 27 |
| | 26.40 | 18 |
| | 27.22 | 14. |

Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

699, 748, 762, 845, 947, 1009, 1035, 1067, 1103, 1118, 1129, 1174, 1193, 1206, 1222, 1247, 1267, 1317, 1365, 1422, 1436, 1456, 1465, 1502, 1592, 1607, 1688, 2412, 2489, 2627, 2846, 2868, 2913, 2928, 3435 cm^{-1} .

(4) Polymorphs (IV)

Peaks in the powder X-ray diffraction pattern are:

| | Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|----|---|--------------------------|
| 40 | 6.24 | 15 |
| | 9.66 | 12 |
| | 11.04 | 22 |
| | 12.12 | 24 |
| 45 | 12.54 | 67 |
| | 12.76 | 61 |
| | 13.98 | 27 |
| | 14.42 | 15 |
| | 14.88 | 11 |
| 50 | 16.34 | 12 |
| | 17.46 | 100 |
| | 18.12 | 25 |
| | 18.60 | 32 |
| 55 | 19.06 | 15 |
| | 19.98 | 74 |
| | 20.42 | 41 |

(continued)

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 20.62 | 34 |
| 21.30 | 48 |
| 21.80 | 63 |
| 22.58 | 78 |
| 23.04 | 46 |
| 24.00 | 32 |
| 24.54 | 49 |
| 25.14 | 90 |
| 25.36 | 99 |
| 26.06 | 34 |
| 28.10 | 41 |
| 28.58 | 39 |
| 29.30 | 31 |
| 29.44 | 28 |

Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

401, 431, 459, 467, 490, 506, 518, 561, 586, 606, 631, 651, 709, 758, 766, 857, 944, 1009, 1041, 1106, 1119, 1132, 1213, 1225, 1265, 1304, 1318, 1429, 1458, 1470, 1500, 1589, 1605, 1630, 1647, 1683, 2562, 2577, 2608, 2634, 2689, 2717, 2836, 2924, 2949, 2989, 3007, 3032, 3061, 3322, 3376, 3422 cm^{-1} .

2. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in Claim 1, which comprises the steps of dissolving Donepezil hydrochloride in ethanol and adding diethyl ether or isopropyl ether to the solution.
3. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in Claim 1, which comprises the steps of dissolving Donepezil hydrochloride in ethanol, adding isopropyl ether to the solution, stirring the mixture for 10 to 30 minutes after having precipitated and filtrating the crystalline precipitates.
4. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in Claim 1, which comprises the steps of dissolving Donepezil and hydrochloric acid or hydrogen chloride in ethanol and adding diethyl ether to the solution.
5. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in Claim 1, which comprises the steps of dissolving Donepezil in ethanol, adding hydrochloric acid or hydrogen chloride to the solution and concentrating the mixture.
6. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in Claim 1, which comprises the steps of dissolving Donepezil in ethanol, adding hydrochloric acid or hydrogen chloride to the solution and adding isopropyl ether to the mixture.
7. The process as claimed in Claim 6, which further comprises filtrating crystalline precipitates after having stirred for 10 to 60 minutes.
8. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 1, which comprises the step of humidifying the polymorph (II) as defined in claim 1.
9. Use of a pharmacologically effective amount of Donepezil hydrochloride in the form of a polymorph as defined in claim 1 for the preparation of a medicament for the treatment of a disease accompanied by acetylcholinesterase activity.
10. Use according to claim 9, in which the disease is senile dementia.

11. Use according to claim 9, in which the disease is senile dementia of the Alzheimer type.
12. A therapeutical composition which comprises a pharmacologically effective amount of Donepezil hydrochloride in the form of polymorph as defined in claim 1 and a pharmacologically acceptable carrier.
13. The Donepezil hydrochloride as claimed in claim 1, which is in the form of polymorph (II).
14. The Donepezil hydrochloride as claimed in claim 1, which is in the form of polymorph (IV).
15. Donepezil hydrochloride, 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride, in the form of polymorphs selected from the group consisting of (II), (IV) and (V), each Polymorphs being specified by peaks at below shown diffraction degrees with the below shown intensity in terms of I/I_0 in powder X-ray diffraction pattern and the below shown absorption peaks in infrared absorption spectra in potassium bromide in terms of reciprocal centimeters:

(2) Polymorphs (II)

Peaks in the powder X-ray diffraction pattern are:

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 10.10 | 76 |
| 12.64 | 14 |
| 15.74 | 85 |
| 15.82 | 86 |
| 16.20 | 100 |
| 16.46 | 87 |
| 17.40 | 50 |
| 17.50 | 48 |
| 17.88 | 31 |
| 18.36 | 28 |
| 18.58 | 51 |
| 18.66 | 46 |
| 19.48 | 42 |
| 20.18 | 81 |
| 20.80 | 36 |
| 22.26 | 45 |
| 23.38 | 86 |
| 23.52 | 59 |
| 24.06 | 34 |
| 24.32 | 55 |
| 25.14 | 44 |
| 25.44 | 50 |
| 25.72 | 39 |
| 25.96 | 35 |
| 26.14 | 25 |
| 28.06 | 25 |
| 28.20 | 34 |
| 28.38 | 34. |

Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

560.1, 698.9, 749.1, 846.2, 947.6, 1036.1, 1119.3, 1222.7, 1266.4, 1318.7, 1364.1, 1458.3, 1500.9, 1522.3, 1534.0, 1542.6, 1560.2, 1570.3, 1592.0, 1637.0, 1647.9, 1654.4, 1689.5, 1718.3, 1734.7, 1751.7, 1773.9, 1793.8, 1830.7, 1846.0, 1870.1, 2345.1, 2489.9, 2927.9, 3448.1 cm^{-1} .

EP 1 211 243 A1

(4) Polymorphs (IV)

Peaks in the powder X-ray diffraction pattern are:

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 9.64 | 11 |
| 10.92 | 11 |
| 12.46 | 63 |
| 12.72 | 17 |
| 13.86 | 27 |
| 14.42 | 12 |
| 17.36 | 100 |
| 18.54 | 39 |
| 19.90 | 37 |
| 21.18 | 35 |
| 21.74 | 39 |
| 22.48 | 60 |
| 22.96 | 36 |
| 24.10 | 17 |
| 25.28 | 70 |
| 28.00 | 27 |
| 28.50 | 27 |

Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

561.5, 709.0, 766.2, 786.3, 804.9, 857.0, 944.3, 979.3, 1041.5, 1118.7, 1264.6, 1318.7, 1364.1, 1458.1, 1499.2, 1542.5, 1560.1, 1588.1, 1636.6, 1647.8, 1654.3, 1684.3, 1718.2, 1734.4, 1751.4, 1773.7, 1793.5, 1830.5, 1845.8, 1870.1, 2344.8, 2369.3, 2719.2, 2922.9, 3324.0 cm^{-1} .

(5) Polymorphs (V)

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 6.58 | 7 |
| 6.86 | 27 |
| 10.12 | 32 |
| 12.54 | 33 |
| 12.90 | 43 |
| 13.64 | 64 |
| 15.58 | 27 |
| 17.22 | 69 |
| 18.44 | 72 |
| 18.96 | 19 |
| 19.30 | 25 |
| 19.64 | 19 |
| 19.74 | 25 |
| 20.30 | 19 |
| 20.46 | 17 |
| 21.10 | 15 |
| 21.96 | 100 |
| 22.24 | 32 |
| 24.22 | 63 |

(continued)

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 24.66 | 96 |
| 25.36 | 60 |
| 26.14 | 15 |
| 26.82 | 44 |
| 27.52 | 24 |
| 27.96 | 15 |
| 28.20 | 49 |
| 29.58 | 13 |
| 29.66 | 17 |
| 29.76 | 17. |

Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

506.5, 559.7, 594.4, 698.0, 740.8, 805.1, 861.9, 948.5, 972.1, 1039.9, 1120.8, 1220.7, 1264.8, 1314.6, 1364.1, 1458.0, 1499.5, 1542.5, 1560.2, 1592.1, 1692.9, 2500.1, 2924.2, 2998.9, 3422.1 cm^{-1} .

16. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil in ethanol, adding hydrochloric acid or hydrogen chloride to the solution and adding tert-butyl methyl ether successively.
17. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil in isopropyl alcohol or in methylene chloride and adding hydrochloric acid or hydrogen chloride to the solution.
18. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil in acetone, adding hydrochloric acid or hydrogen chloride to the solution and filtrating the crystalline precipitates within 30 minutes from the separation of crystals.
19. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil hydrochloride in ethanol and adding tert-butyl methyl ether to the solution at temperature in the range of 10 to 30°C.
20. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil hydrochloride in ethanol and adding this solution to diisopropyl ether.
21. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil hydrochloride in methylene chloride and adding tert-butyl methyl ether to the solution.
22. A process for producing the polymorph (II) of Donepezil hydrochloride as defined in claim 15, which comprises the step of drying the polymorph (I) or amorphous form of Donepezil hydrochloride, wherein polymorph (I) is defined as follows:

(1) Polymorphs (I)

Peaks in the powder X-ray diffraction pattern are:

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 9.82 | 13 |
| 10.48 | 18 |
| 12.52 | 93 |
| 13.02 | 69 |

(continued)

| Diffraction angles ($2\theta, ^\circ$) | Intensity (I/I_0) |
|---|--------------------------|
| 13.52 | 34 |
| 13.74 | 37 |
| 14.78 | 31 |
| 16.00 | 45 |
| 16.76 | 37 |
| 17.46 | 34 |
| 19.18 | 26 |
| 19.66 | 32 |
| 21.04 | 100 |
| 21.16 | 82 |
| 22.92 | 52 |
| 23.82 | 72 |
| 24.14 | 32 |

Wave numbers (cm^{-1}) of infrared absorption spectra in potassium bromide are:

562.7, 603.2, 700.4, 749.6, 798.1, 859.2, 896.0, 921.3, 946.3, 971.8, 1038.0, 1119.3, 1216.8, 1266.0, 1315.4, 1367.7, 1454.1, 1501.5, 1537.8, 1555.9, 1590.7, 1643.7, 1681.9, 2350.9, 2534.0, 2922.1, 3381.8, 3585.2 cm^{-1} .

23. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in Claim 15, which comprises the step of humidifying the polymorph (II) as defined in Claim 15.
24. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil in water with or without tetrahydrofuran and adding hydrochloric acid or hydrogen chloride to the solution.
25. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil in hydrochloric acid and adding tetrahydrofuran to the solution.
26. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil in toluene and adding hydrochloric acid to the solution.
27. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 15, which comprises the steps of dissolving Donepezil in n-hexane and adding hydrochloric acid to the solution.
28. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 15, which comprises the step of crystallizing Donepezil in a mixture of methanol and hydrochloric acid.
29. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 15, which comprises the step of crystallizing Donepezil hydrochloride from water.
30. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 15, which comprises the step of humidifying the amorphous form of Donepezil hydrochloride.
31. A process for producing the polymorph (IV) of Donepezil hydrochloride as defined in claim 15, which comprises the step of humidifying the polymorph (II) of Donepezil hydrochloride.
32. A process for producing the polymorph (V) of Donepezil hydrochloride as defined in claim 15, which comprises the step of drying the polymorph (IV) of Donepezil hydrochloride.
33. Use of a pharmacologically effective amount of the Donepezil hydrochloride in the form of a polymorph as defined

in claim 15 for the preparation of a medicament for the treatment of a disease accompanied by acetylcholinesterase activity.

34. Use as claimed in claim 33, in which the disease is senile dementia.

35. Use as claimed in claim 33, in which the disease is senile dementia of the Alzheimer type.

36. A therapeutical composition which comprises a pharmacologically effective amount of Donepezil hydrochloride in the form of polymorph as defined in claim 15 and a pharmacologically acceptable carrier.

37. The Donepezil hydrochloride as claimed in claim 15, which is in the form of polymorph (II).

38. The Donepezil hydrochloride as claimed in claim 15, which is in the form of polymorph (IV).

39. The Donepezil hydrochloride as claimed in claim 15, which is in the form of polymorph (V).

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21




Fig. 22




Fig. 23




Fig. 24.




Fig. 25




Fig. 26




Fig. 27




Fig. 28




Fig. 29




Fig. 30




Fig. 31




Fig. 32




Fig. 33




Fig. 34




Fig. 35




Fig. 36




Fig. 37





Fig. 38





European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 02 00 5248

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|---|--|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int.CI.7) |
| D, A | EP 0 296 560 A (EISAI CO., LTD., JAPAN) 28 December 1988 (1988-12-28) * example 4 * | 1-39 | C07D211/32 A61K31/445 A61P25/28 |
| A | EP 0 673 927 A (EISAI CO., LTD.) 27 September 1995 (1995-09-27) * example 4 * | 1-39 | |
| | | | TECHNICAL FIELDS SEARCHED (Int.CI.7) |
| | | | C07D A61K A61P |
| The present search report has been drawn up for all claims | | | |
| Place of search THE HAGUE | | Date of completion of the search 9 April 2002 | Examiner Chouly, J |
| CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document | | | |

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 00 5248

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

09-04-2002

| Patent document cited in search report | | Publication date | Patent family member(s) | Publication date |
|---|---|---------------------|----------------------------|---------------------|
| EP 0296560 | A | 28-12-1988 | AT 134618 T | 15-03-1996 |
| | | | AT 171161 T | 15-10-1998 |
| | | | AT 205828 T | 15-10-2001 |
| | | | AT 204862 T | 15-09-2001 |
| | | | AU 627151 B2 | 20-08-1992 |
| | | | AU 1821688 A | 22-12-1988 |
| | | | CA 1338808 A1 | 24-12-1996 |
| | | | CA 1340192 A1 | 15-12-1998 |
| | | | CN 1030752 A ,B | 01-02-1989 |
| | | | CN 1073939 A ,B | 07-07-1993 |
| | | | CN 1071417 A ,B | 28-04-1993 |
| | | | CY 1940 A | 16-05-1997 |
| | | | DD 283377 A5 | 10-10-1990 |
| | | | DE 3855028 D1 | 04-04-1996 |
| | | | DE 3856251 D1 | 22-10-1998 |
| | | | DE 3856251 T2 | 18-03-1999 |
| | | | DE 3856487 D1 | 04-10-2001 |
| | | | DE 3856491 D1 | 25-10-2001 |
| | | | DK 108296 A | 03-10-1996 |
| | | | DK 108396 A | 03-10-1996 |
| | | | DK 337988 A | 23-12-1988 |
| | | | EP 1116716 A1 | 18-07-2001 |
| | | | EP 0296560 A2 | 28-12-1988 |
| | | | EP 0579263 A1 | 19-01-1994 |
| | | | EP 0673927 A1 | 27-09-1995 |
| | | | EP 0742207 A1 | 13-11-1996 |
| | | | ES 2083359 T3 | 16-04-1996 |
| | | | ES 2121039 T3 | 16-11-1998 |
| | | | ES 2164720 T3 | 01-03-2002 |
| | | | ES 2160747 T3 | 16-11-2001 |
| | | | FI 882716 A ,B, | 23-12-1988 |
| | | | FI 952850 A | 09-06-1995 |
| | | | FI 962753 A | 04-07-1996 |
| | | | GR 3019142 T3 | 31-05-1996 |
| | | | HK 216396 A | 27-12-1996 |
| | | | HU 50768 A2 | 28-03-1990 |
| | | | HU 9500122 A3 | 28-07-1995 |
| | | | HU 9500123 A3 | 28-06-1995 |
| | | | JP 1079151 A | 24-03-1989 |
| | | | JP 2578475 B2 | 05-02-1997 |
| | | | JP 3078244 B2 | 21-08-2000 |
| | | | JP 10067739 A | 10-03-1998 |
| | | | JP 2733203 B2 | 30-03-1998 |
| | | | JP 7252216 A | 03-10-1995 |
| | | | KR 9103618 B1 | 07-06-1991 |
| | | | LU 90221 A9 | 06-05-1998 |

EPO FORM P445

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 00 5248

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

09-04-2002

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| EP 0296560 A | | NO 177590 B | 10-07-1995 |
| | | NZ 224970 A | 25-09-1991 |
| | | PH 26315 A | 29-04-1992 |
| | | PT 87783 A ,B | 01-07-1988 |
| EP 673927 A | 27-09-1995 | EP 1116716 A1 | 18-07-2001 |
| | | EP 0579263 A1 | 19-01-1994 |
| | | EP 0673927 A1 | 27-09-1995 |
| | | EP 0742207 A1 | 13-11-1996 |
| | | AT 134618 T | 15-03-1996 |
| | | AT 171161 T | 15-10-1998 |
| | | AT 205828 T | 15-10-2001 |
| | | AT 204862 T | 15-09-2001 |
| | | AU 627151 B2 | 20-08-1992 |
| | | AU 1821688 A | 22-12-1988 |
| | | CA 1338808 A1 | 24-12-1996 |
| | | CA 1340192 A1 | 15-12-1998 |
| | | CN 1030752 A ,B | 01-02-1989 |
| | | CN 1073939 A ,B | 07-07-1993 |
| | | CN 1071417 A ,B | 28-04-1993 |
| | | CY 1940 A | 16-05-1997 |
| | | DD 283377 A5 | 10-10-1990 |
| | | DE 3855028 D1 | 04-04-1996 |
| | | DE 3856251 D1 | 22-10-1998 |
| | | DE 3856251 T2 | 18-03-1999 |
| | | DE 3856487 D1 | 04-10-2001 |
| | | DE 3856491 D1 | 25-10-2001 |
| | | DK 108296 A | 03-10-1996 |
| | | DK 108396 A | 03-10-1996 |
| | | DK 337988 A | 23-12-1988 |
| | | EP 0296560 A2 | 28-12-1988 |
| | | ES 2083359 T3 | 16-04-1996 |
| | | ES 2121039 T3 | 16-11-1998 |
| | | ES 2164720 T3 | 01-03-2002 |
| | | ES 2160747 T3 | 16-11-2001 |
| | | FI 882716 A ,B, | 23-12-1988 |
| | | FI 952850 A | 09-06-1995 |
| | | FI 962753 A | 04-07-1996 |
| | | GR 3019142 T3 | 31-05-1996 |
| | | HK 216396 A | 27-12-1996 |
| | | HU 50768 A2 | 28-03-1990 |
| | | HU 9500122 A3 | 28-07-1995 |
| | | HU 9500123 A3 | 28-06-1995 |
| | | JP 1079151 A | 24-03-1989 |
| | | JP 2578475 B2 | 05-02-1997 |
| | | JP 3078244 B2 | 21-08-2000 |

EPO FORM P/USE

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 00 5248

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

09-04-2002

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| EP 673927 A | JP | 10067739 A | 10-03-1998 |
| | JP | 2733203 B2 | 30-03-1998 |
| | JP | 7252216 A | 03-10-1995 |
| | KR | 9103618 B1 | 07-06-1991 |
| | LU | 90221 A9 | 06-05-1998 |
| | NO | 177590 B | 10-07-1995 |
| | NZ | 224970 A | 25-09-1991 |
| | PH | 26315 A | 29-04-1992 |
| | PT | 87783 A , B | 01-07-1988 |
| <hr/> | | | |

EPO FORM P4455

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)